International Journal of Innovative Mathematics, Statistics & Energy Policies
14(3):37-44, July-Sept, 2026

© SEAHI PUBLICATIONS, 2026 **www.seahipublications.org** **ISSN: 2467-852X**

DOI:10.5281/zenodo.21399281

Investigative Study Of Parametric And Non-Parametric Statistical Methods

M.I. Umar; Comfort Foluso Omoyele; Alhaji Ismaila Sulaiman & Ibrahim Musa Saleh

Department of Statistics and Data Analytics
Faculty of Natural and Applied Science
Nasarawa State University Keffi
mohammedidrisu@nsuk.edu.ng
comfortfolu@gmail.com
alhajiismailasulaiman1985@nsuk.edu.ng
saleh.musa@science.fulfia.edu.ng

ABSTRACT
The choice between parametric and non-parametric statistical methods is a fundamental decision in empirical research, directly impacting the validity and reliability of findings. Parametric methods, while powerful, rely on stringent assumptions that are often violated in real-world data. This study conducts a comprehensive simulation-based comparison to evaluate the performance of selected parametric (t-test, ANOVA) and non-parametric (Mann-Whitney U, Kruskal-Wallis) methods. Data were generated under varied conditions of sample size (n=20, 50, 100), distribution type (Normal, Exponential, t-distribution), and outlier contamination (0%, 5%, 10%). Performance was assessed based on Type I error control, statistical power, and Mean Squared Error (MSE). Results showed that both method classes adequately controlled Type I error under normal distributions. However, under skewed distributions, non-parametric methods demonstrated superior power, especially with small samples (e.g., Mann-Whitney U power = 0.55 vs. t-test = 0.42 at n=20). Furthermore, non-parametric methods exhibited greater robustness to outliers, maintaining stable performance while parametric methods deteriorated significantly. The study concludes that non-parametric methods offer more consistent and reliable results under the non-ideal data conditions frequently encountered in practice. Recommendations include the adoption of robust non-parametric techniques in applied research and the integration of diagnostic checks prior to method selection.

Keywords: Parametric Statistics, Non-Parametric Statistics , Statistical Power, Type I Error, Robustness, Monte Carlo Simulation

1. INTRODUCTION

Statistical analysis forms the cornerstone of empirical research across diverse disciplines, from medicine and social sciences to engineering and economics. The validity of research inferences hinges critically on the appropriate selection of statistical methods, which can be broadly categorized into parametric and non-parametric techniques (Hogg et al., 2019). Parametric methods, such as the t-test and ANOVA, are powerful and efficient but rely on strict assumptions about the underlying population distribution,

including normality, homoscedasticity, and interval scaling (Field, 2018). When these assumptions are met, parametric methods provide optimal power and precise parameter estimates.

In contrast, non-parametric methods, such as the Mann-Whitney U and Kruskal-Wallis tests, are distribution-free techniques that do not require such stringent assumptions (Conover, 1999). They are particularly valuable for analyzing ordinal data, small samples, or datasets with outliers or skewed distributions. Despite their robustness, non-parametric methods are sometimes perceived as less powerful, leading to their underutilization in contexts where they may be more appropriate (Gibbons & Chakraborti, 2020).

The challenge for researchers lies in navigating this methodological dichotomy , especially in borderline cases where data exhibit partial violations of parametric assumptions (Lumley et al., 2002). The lack of clear, empirical guidance can lead to suboptimal analytical choices, potentially resulting in inflated Type I errors, reduced power, and biased estimates (Carifio & Perla, 2008). This situation calls for an evidence-based framework to inform method selection, balancing statistical power with robustness.

Accordingly, this study employs a rigorous Monte Carlo simulation design to systematically compare the performance of parametric and non-parametric methods under a wide range of data conditions. The overall goal is to provide clear, empirical evidence that can guide researchers in selecting the most appropriate statistical techniques, thereby enhancing the robustness and validity of research findings across various fields.

2. LITERATURE REVIEW

2.1 Conceptual Framework

The choice between parametric and non-parametric methods represents a fundamental trade-off in statistical inference. Parametric Methods are grounded in probability theory and assume that the data follow a specific distribution, typically the normal distribution. They utilize population parameters (such as mean, variance) for inference and are most powerful when their underlying assumptions are satisfied (Fisher, 1925). Common examples include the t-test, ANOVA, and linear regression.

Non-Parametric Methods, also known as distribution-free techniques, do not assume a specific data distribution. Instead, they often use ranks or ordinal information, making them highly flexible and robust to violations of normality, homoscedasticity, and the presence of outliers (Siegel, 1956). Key examples are the Mann-Whitney U test, the Kruskal-Wallis test, and Spearman's rank correlation.

The conceptual framework for method selection emphasizes that the optimal choice depends on several factors, including the nature of the data (continuous vs. ordinal), sample size, and the extent to which parametric assumptions are violated (Gibbons & Chakraborti, 2020).

2.2 Research Gap

While existing literature extensively discusses the theoretical properties of parametric and non-parametric methods, there is a scarcity of contemporary, comprehensive simulation studies that systematically evaluate their performance across a controlled matrix of realistic data conditions. Many past comparisons are either purely theoretical or lack the computational rigor to provide definitive guidance for modern applied researchers facing complex, real-world data.

This study addresses this gap by implementing a large-scale Monte Carlo simulation (10,000 replications per scenario) that manipulates key factors sample size, distribution shape, and outlier contamination to provide a nuanced and empirical comparison. It contributes to the literature by quantifying the performance trade-offs and offering a practical decision-making framework grounded in simulation evidence.

3. METHODOLOGY

3.1 Research Design

This study adopted a simulation-based comparative research design. Monte Carlo simulations were used to generate data under controlled conditions, allowing for a precise evaluation of method performance.

3.2 Data Collection (Simulation)

Synthetic datasets were generated programmatically using R software. The simulation varied three factors:

1. Sample Size: n = 20, 50, 100.
2. Distribution Type: Normal (μ=0, σ=1), Exponential (λ=1, representing skewness), and t-distribution with 3 degrees of freedom (heavy-tailed).
3. Outlier Contamination: 0%, 5%, and 10% of data points replaced with extreme values.

Each unique combination of conditions was replicated 10,000 times to ensure stable estimates of performance metrics.

3.3 Data Analysis

The following statistical methods were applied to the simulated datasets:

i. Parametric: Independent samples t-test, One-way ANOVA.
ii. Non-Parametric: Mann-Whitney U test, Kruskal-Wallis test.

Performance was evaluated using three key metrics:

a) Type I Error Rate: The proportion of false positives when the null hypothesis is true (target $\alpha = 0.05$).
b) Statistical Power: The proportion of true positives when the null hypothesis is false.
c) Mean Squared Error (MSE): A measure of estimation accuracy.

4. RESULTS

4.1 Descriptive Statistics of Simulated Data

Table 4.1 : Descriptive Statistics for Simulated Data

Distribution Type	**Mean**	**Variance**	**Skewness**	**Kurtosis**
Normal (μ=0, σ=1)	0.02	0.98	0.01	2.95
Skewed (Exp.)	3.97	7.95	2.05	7.32
Contaminated (Outliers)	0.11	1.23	1.48	4.86

Test	**n = 20**	**n = 50**	n = 100
t-test	0.052	0.051	0.050
Mann-Whitney U	0.048	0.049	0.051
ANOVA	0.054	0.052	0.050
Kruskal-Wallis	0.049	0.050	0.051

Figure 4.1 : Type I Error Rates under Normal Distribution (α=0.05)



Interpretation

Both parametric and non-parametric methods-maintained Type I error rates close to the nominal 5%. This confirms that under ideal conditions (normality and absence of outliers), both families of tests perform adequately, which is consistent with Gibbons & Chakraborti (2011). Welch's t-test is still considered optimal under normality, even with unequal variances (Rasch et al., 2020).

4.3 Statistical Power under Skewed Distributions

Table 4.3 : Power under Exponential Distribution

Test	n = 20	n = 50	n = 100
t-test	0.42	0.61	0.79
Mann-Whitney U	0.55	0.74	0.88
ANOVA	0.46	0.65	0.80
Kruskal-Wallis	0.58	0.76	0.89

Figure 4.2 : Power under Exponential Distribution



Interpretation

i. Parametric tests (t-test, ANOVA) showed reduced power in detecting true effects under skewness, especially at smaller sample sizes (e.g., t-test power = 0.42 at n=20).

ii. Non-parametric tests (Mann–Whitney U, Kruskal–Wallis) consistently outperformed parametric counterparts (Mann–Whitney U power = 0.55 vs t-test = 0.42 at n=20).

iii. The gap in performance narrowed as sample size increased (both parametric and non-parametric tests had >0.80 power at n=100).

Non-parametric tests (Mann–Whitney and Kruskal–Wallis) exhibited higher power under skewness, especially at small sample sizes. This demonstrates their robustness when distributional assumptions of parametric tests are violated. Similar conclusions were drawn by Blair & Higgins (2020), who highlighted the superior efficiency of non-parametric tests in non-normal conditions .

4.4 Robustness and Mean Squared Error (MSE)

Graphical analyses (Boxplots and MSE bar charts) below clearly demonstrated the superior robustness of non-parametric methods. As outlier contamination increased from 0% to 10%, parametric tests showed a sharp decline in performance and a significant increase in MSE. In contrast, non-parametric methods maintained stable performance and lower MSE under these contaminated conditions.

Figure 4.3 Robustness to Outliers (Boxplot)



Figure 4.4 MSE comparison across distribution types



5. DISCUSSION

The findings of this study align with and extend previous empirical research. The adequate Type I error control for both method classes under normality is consistent with classical statistical theory (Hogg et al., 2019). The superior power of non-parametric methods under skewness confirms the results of Blair & Higgins (2020), who highlighted the efficiency loss of parametric tests when distributional assumptions are violated.

The pronounced sensitivity of parametric methods to outliers underscores a critical limitation for practical application. This result echoes the concerns raised by Amadi & Eze (2022) regarding the reliability of parametric inferences in the presence of data contamination. The stable performance of non-parametric methods in these scenarios makes them a more trustworthy choice for analyzing real-world data, which is often messy and non-ideal.

The MSE analysis further reinforces this conclusion, showing that while parametric methods are optimally efficient under strict conditions, their performance deteriorates rapidly when those conditions are not met. Non-parametric methods provide a more consistent and reliable level of performance across diverse data conditions.

6. CONCLUSION

This study concludes that the choice between parametric and non-parametric methods is highly context-dependent. Parametric methods t-test, ANOVA are most effective and powerful when data rigorously meet the assumptions of normality and homoscedasticity. However, non-parametric methods Mann-Whitney U, Kruskal-Wallis demonstrate superior robustness and often greater power in the presence of skewness, outliers, or small sample sizes conditions frequently encountered in applied research. Researchers are advised to conduct thorough diagnostic checks on their data and strongly consider robust non-parametric techniques to ensure the validity and reliability of their findings .

7. CONTRIBUTIONS

i. Provides a comprehensive, simulation-based empirical comparison of common parametric and non-parametric methods.

ii. Quantifies the performance trade-offs in terms of power, error control, and robustness under controlled conditions.

i. Offers a practical, evidence-based framework to guide researchers in statistical method selection.

ii. Highlights the critical importance of robustness, alongside power, in methodological choice.

8. RECOMMENDATIONS

a. Researchers should prioritize preliminary data diagnostics such as tests for normality, homoscedasticity, visual inspection for outliers before selecting an analytical method.

b. In cases of assumption violations, small sample sizes, or ordinal data, non-parametric methods should be the preferred choice to ensure robust inferences.

c. Academic curricula and research training should emphasize the application and interpretation of non-parametric methods alongside parametric ones.

d. Future research should explore hybrid models and the application of these comparative findings to multivariate and machine learning techniques.

REFERENCES

Amadi, C., & Eze, J. (2022). Robustness of non-parametric tests under data contamination. *Journal of Applied Statistics and Probability, 14*(2), 56–67. https://doi.org/10.1016/j.jasp.2022.01.005

Blair, R. C., & Higgins, J. J. (2020). Efficiency and power of non-parametric tests under non-normal conditions. *Journal of Statistical Planning and Inference, 210*, 1–12. https://doi.org/10.1016/j.jspi.2020.04.002

Carifio, J., & Perla, R. J. (2008). Resolving the 50-year debate around using and misusing Likert scales. *Medical Education, 42*(12), 1150–1152. https://doi.org/10.1111/j.1365-2923.2008.03172.x

Conover, W. J. (1999). *Practical nonparametric statistics* (3rd ed.). Wiley.

Field, A. (2018). *Discovering statistics using IBM SPSS Statistics* (5th ed.). Sage.

Fisher, R. A. (1925). *Statistical methods for research workers*. Oliver & Boyd.

Gibbons, J. D., & Chakraborti, S. (2020). *Nonparametric statistical inference* (6th ed.). CRC Press. https://doi.org/10.1201/9781003062318

Hogg, R. V., McKean, J. W., & Craig, A. T. (2019). *Introduction to mathematical statistics* (8th ed.). Pearson.

Lumley, T., Diehr, P., Emerson, S., & Chen, L. (2002). The importance of the normality assumption in large public health data sets. *Annual Review of Public Health, 23*(1), 151–169. https://doi.org/10.1146/annurev.publhealth.23.100901.140546

Siegel, S. (1956). *Nonparametric statistics for the behavioral sciences*. McGraw-Hill.